

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11)

EP 0 792 292 B1

(12)

# EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention  
of the grant of the patent:  
18.08.1999 Bulletin 1999/33

(21) Application number: 96931047.3

(22) Date of filing: 09.09.1996

(51) Int Cl.<sup>6</sup>: C07K 14/705

(86) International application number:  
PCT/EP96/03933

(87) International publication number:  
WO 97/09348 (13.03.1997 Gazette 1997/12)

(54) ORPHAN RECEPTOR  
WAISEN-REZEPTOR  
RECEPTEUR ORPHELIN

(84) Designated Contracting States:  
AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC  
NL PT SE

(30) Priority: 08.09.1995 GB 9518272  
15.03.1996 GB 9605550  
11.04.1996 GB 9607532  
08.05.1996 GB 9609576

(43) Date of publication of application:  
03.09.1997 Bulletin 1997/36

(60) Divisional application: 99200009.1 / 0 935 000

(73) Proprietor: KARO BIO AB  
141 57 Huddinge (SE)

(72) Inventors:  
• KUIPER, George G.J., M.-Karolinska Institute  
Novum, 141 86 Huddinge (SE)  
• ENMARK, Eva - Karolinska Institute  
Novum, 141 96 Huddinge (SE)  
• GUSTAFSSON, Jan-Ake - Karolinska Institute  
Novum, 141 86 Huddinge (SE)

(74) Representative: Dean, John Paul et al  
Withers & Rogers,  
Goldings House,  
2 Hays Lane  
London SE1 2HW (GB)

(56) References cited:

- NUCLEIC ACIDS RESEARCH, vol. 15, no. 6, 25 March 1987, pages 2499-2513, XP002026307 KOIKE, S. ET AL.: "Molecular cloning and characterization of a rat estrogen receptor cDNA"
- SCIENCE, vol. 231, no. 4742, 7 March 1986, pages 1150-1154, XP000611679 GREENE G L ET AL.: "SEQUENCE AND EXPRESSION OF HUMAN ESTROGEN RECEPTOR COMPLEMENTARY DNA"
- NUCLEIC ACIDS RESEARCH, vol. 17, no. 14, 25 July 1989, pages 5477-5488, XP002026308 LEES J. ET AL.: "Identification of two transactivation domains in the mouse oestrogen receptor"
- PNAS U.S.A., vol. 93, no. 12, 11 June 1996, pages 5925-5930, XP002026309 KUIPER G. ET AL.: "Cloning of a novel estrogen receptor expressed in rat prostate and ovary"
- FEBS LETTERS, vol. 392, no. 1, 19 August 1996, pages 49-53, XP002026310 MOSSELMAN, S. ET AL.: "ERbeta: identification and characterization of a novel human estrogen receptor"
- MOLECULAR ENDOCRINOLOGY, vol. 10, no. 11, November 1996, pages 1293-1307, XP000645166 ENMARK E. AND GUSTAFSSON J.: "Orphan nuclear receptors- the first eight years"

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 0 792 292 B1



## Description

[0001] This invention relates to cellular nuclear receptors and their uses..

[0002] A large family of nuclear receptors which confer cells with responsiveness to molecules such as retinoid acid, vitamin D, steroid hormones and thyroid hormones has been identified. Extensive studies have shown that the members of this superfamily of nuclear receptors activate and/or repress gene transcription through direct binding to discrete cis-acting elements termed "hormone response elements" (HRE). It has been shown that these HRE's comprise repeats of consensus palindromic hexanucleotide DNA motifs. The specificity of the HRE's is determined by the orientation of, and spacing between, halfsites (i.e. half a palindromic sequence) (Umenesono K., *et al*, 1991 *Cell* 65, 1255-1266).

[0003] Specific DNA binding is mediated by a strongly-conserved DNA binding domain, containing two zinc fingers, which is conserved among all thus discovered nuclear receptors. Three amino acids at the C-terminal base of the first zinc finger (known as the "P-box") are important for the recognition of the half site nucleotide sequence. Members of the nuclear receptor superfamily have been classified into different groups on the basis of the amino acid sequence within the P box.

[0004] All members of the nuclear receptor superfamily also contain a hypervariable N-terminal domain and a ligand-binding domain containing some "patches" of conserved sequence. One of these is called the "Ti-domain".

[0005] Molecules which are thought to be nuclear receptors, as they are structurally related to characterised receptors, but for which no ligand has been found, are termed "orphan receptors". Many such orphan receptors have been identified (see for example Evans R.M. (1988) *Science* 240, 889-895 and O'Malley, B. (1990) *Mol. Endocrinol.* 4 363-369).

[0006] We have now unexpectedly identified, initially in rat a new orphan receptor, which is related to the known estrogen receptor ER $\alpha$ , and which we have designated "ER $\beta$ " (specifically "rER $\beta$ " in rat). In this specification "Er $\beta$ " will be used to refer to the receptors hER $\beta$  or rER $\beta$  or related receptors. The nucleotide and amino acid sequences of rER $\beta$  have now been determined and are shown in Fig. 1. We have also identified a human Er $\beta$  - "hER $\beta$ ", the amino acid DNA and sequences of which are shown in Fig. 13A and 13B respectively.

[0007] According to one aspect of the invention there is provided an estrogen receptor (called ER $\beta$ ) having the amino acid sequence of Fig. 1, and an estrogen receptor having an amino acid sequence which is more than 95% identical with that sequence; an estrogen receptor (called ER $\beta$ ) having the amino acid sequence of Fig. 13A, and an estrogen receptor which is more than 89% identical with that amino acid sequence, and which is relatively highly expressed in HUVEC and HOS D4 cells compared to the expression of ER $\alpha$  in those cells and an estrogen receptor (called ER $\beta$ ) having the amino acid sequence of Fig. 14A and an estrogen receptor having an amino acid sequence which is more than 95% identical with that amino acid sequence. The isolated receptor may be particularly useful in the search for molecules for use in treatment of diseases or conditions such as cardiovascular diseases, central nervous system diseases or conditions or osteoporosis, prostate cancer or benign prostatic hyperplasia.

[0008] The receptor of the invention may also be used in the testing of environmental chemicals for estrogenic activity. There has been increasing concern over the effect of various chemicals released into the environment on the reproduction of humans and animals. Threats to the reproductive capabilities of birds, fish, reptiles, and some mammals have become evident and similar effects in humans have been proposed. Substantial evidence is now emerging which shows that exposure to certain chemicals during critical periods of foetal life may distort the development of the reproductive organs and the immune and nervous systems. On the basis of possible parallels between actual wildlife effects, seen for example in birds and seals living in highly polluted areas, and proposed effects in humans, in combination with documented human reproductive effects caused by prenatal exposure to the pharmaceutical estrogen, diethyl stilbestrol (DES), "estrogenic" chemicals have been proposed to threaten the reproductive capability of both animals and humans. Among the chemicals known or suspected to act as estrogen mimics on the human body, or in other ways disturb the human endocrine system, there are several which have already been identified as environmental hazards. Among the chemicals that have been mentioned as potential causes of disruption of reproductive function in animals and humans are chlorinated organic compounds such as dieldrin, endosulfans, chlordanes, endrins, aldrin, DDT and some PCBs, plastics such as Bisphenol A, phthalates and nonylphenol, and aromatic hydrocarbons. Some of the proposed effects on humans have been suggested to be due to an increasing exposure to environmental estrogens - in fact, exposure to chemical compounds to which higher organisms during the foetal period react in a way that is similar to when they are exposed to high dosages of estrogens. The effects are manifested by for example perturbations of the sex characteristics and impaired reproductive potential. In humans, elevated risks of breast cancer and other hormone-related disease has also been discussed as possible effects. In addition, to the documented "estrogenic" effects, it has recently been demonstrated that environmental pollutants may also act on hormonal pathways other than the estrogenic pathway - it has been shown that p,p' - DDE the main metabolite of DDT (also in humans) is a fairly anti-androgenic agent (Kelce W.R. *et al* *Nature* 1995 375:581-585). Epidemiological studies on these issues are, however, presently difficult to interpret. Nevertheless, there is a growing opinion against these potentially hormone disrupting chemicals, and very palpable public and environmental demand for the governmental agencies and industry



to act. In view of the similarities between the receptor of the present invention, Er $\beta$  and the classical estrogen receptor, Er $\beta$  may be used in the testing of chemicals for estrogenic effect.

[0009] According to another aspect of the invention there is provided a DNA sequence encoding an estrogen receptor according to the first aspect of the invention.

[0010] Er $\beta$  is unique in that it is extremely homologous to the rat estrogen receptor, in particular in its DNA binding domain. It appears that ER $\beta$  has a very limited tissue distribution. In female rats, it appears to be present only in the ovaries, and in male rats in the prostate and testes. As these tissues are classic targets for estrogen action, it can be deduced that ER $\beta$  may mediate some of the effects of estrogen.

[0011] The different ligand specificity of ER $\alpha$  and ER $\beta$  may be exploited to design pharmaceutical agents which are selective for either receptor. In particular, the differences in ligand specificity may be used to develop drugs that specifically target cardiovascular disease in postmenopausal women or osteoporosis.

[0012] The nuclear receptor of the invention, ER $\beta$ , a method of producing it, and tests on its functionality will now be described, by way of example only, with reference to the accompanying drawings, Figs. 1 to 15 in which:

Fig. 1 shows the amino acid sequence of ER $\beta$  and the nucleotide sequence of the gene encoding it;  
 Fig. 2A is a phylogenetic tree showing the evolution of ER $\beta$  and other receptors;  
 Fig. 2B shows the homology between the different domains in ER $\beta$  and certain other receptors;  
 Fig. 2C is an alignment of the amino acid sequence in the ligand binding domains of rER $\beta$ , rER $\alpha$ , mER $\alpha$  and hER $\alpha$ ;  
 Fig. 2D is an alignment of the amino acid sequence in the DNA binding domains of rER $\beta$ , rER $\alpha$ , mER $\alpha$  and hER $\alpha$ ;  
 Fig. 3A is a film autoradiograph of prostate gland showing strong expression of a clone of the receptor of the invention, clone 29;  
 Fig. 3B is a darkfield image showing prominent signal for clone 29 in epithelium (e) of prostatic alveoli. The stroma (s) exhibit(s) weaker signal;  
 Fig. 3C is a bipolarization image of cresyl violet counterstained section showing silver grains over epithelium (e), whereas the stroma(s) contain(s) less grains;  
 The bar represents 0.7 mm for Fig. 3A, 200  $\mu$ m for Fig. 3B and 30  $\mu$ m for Fig. 3C;  
 Fig. 4A shows a film autoradiograph of ovary showing strong expression of clone 29 in follicles at different developmental stages (some are indicated by arrows). The interstitial tissue (arrowheads) shows low signal;  
 Fig. 4B shows a darkfield image showing high expression of clone 29 in granular cells of primary (1), secondary (2), tertiary (3) and mature (4) follicles. Low signal is present in interstitial tissue (it);  
 Fig. 4C is a bipolarization image of ovary showing strong signal in granular cells (gc), whereas the oocyte (oc) and the caintema (ti) are devoid of clear signal;  
 The bar represents 0.9 mm for Fig. 4A, 140  $\mu$ m for Fig. 4B and 50  $\mu$ m for Fig. 4C;  
 Fig. 5A illustrates the results of saturation ligand binding analysis of cloned ER $\beta$ ;  
 Fig. 5B illustrates the specificity of ligand binding by cloned ER $\beta$ ;  
 Fig. 5C illustrates E2 binding by ER $\beta$ ;  
 Fig. 6 illustrates the activation of transcription by cloned ER $\beta$ ;  
 Fig. 7 and 7A illustrates stimulation by various ligands by cloned ER $\beta$ ;  
 Fig. 8 illustrates the results of RT-PCR experiments on the expression of rat estrogen receptors;  
 Fig. 9 illustrates the results of RT-PCR experiments on the expression of human Er $\beta$  (hER $\beta$ );  
 Fig. 10A is a Hill plot comparing binding of <sup>125</sup>I-E2 by hER $\alpha$  and rER $\beta$ ;  
 Fig. 10B is a Scatchard plot comparing binding of <sup>125</sup>I-E2 by hER $\alpha$  and rER $\beta$ ;  
 Fig. 11A illustrates the relative binding affinity of hER $\alpha$  and rER $\beta$  for various ligands;  
 Fig. 11B is a detail of Fig. 12A;  
 Fig. 12 is an alignment of various estrogen receptors;  
 Fig. 13A shows the amino acid sequence of human ER $\beta$ ;  
 Fig. 13B shows the DNA sequence of human ER $\beta$ ;  
 Fig. 14A shows the amino acid sequence of mER $\beta$ ;  
 Fig. 14B shows the DNA sequence of mouse ER $\beta$ ; and  
 Fig. 15 illustrates ligand binding affinities for various phytoestrogens by ER's of the invention.

## A. CLONING OF RAT ER $\beta$

### 1. PCR-amplification and complementary DNA cloning.

[0013] A set of degenerate primers (DBD 1.2.3 and WAK/FAK) were designed previously according to the most highly conserved sequences of the DNA-binding domain (P-box) and ligand binding domain (Ti-stretch) of members of the nuclear receptor family (Enmark, E., Kainu, T., Peltto-Huikko, M., & Gustafsson, J-A (1994) *Biochem. Biophys. Res.*



*Commun.* 204, 49-56). Single strand complementary DNA reverse transcribed from rat prostate total RNA was employed with the primers in PCR reactions as described in Enmark, E., Kainu, T., Peltö-Huikko, M., & Gustafsson, J.-Å (1994) *Biochem. Biophys. Res. Commun.* 204, 49-56. The amplification products were separated on a 2% low melting agarose gel and DNA products between 400 and 700 bp were isolated from the gel and ligated to TA cloning vector (Invitrogen). As alternatives, we also used the RP-1/RP-2 and DBD66-100/DBD210-238 primer sets in the DNA-binding domain of nuclear receptors exactly as described by Hirose T., Fijimoto, W., Yamaai, T., Kim, K.H., Matsuura, H., & Jetten, A.M (1994) *Mol. Endocrinol.* 8, 1667-1677 and Chang, C., Lopes Da Silva, S., Ideta, R., Lee, Y., Yeh, S., & Burbach, J.P.H (1994) *Proc. Natl. Acad. Sci.* 91, 6040-6044 respectively. Clone number 29 (obtained with the DBD-WAK/FAK set) with a length of 462 bp showed high homology (65%) with the rat estrogen receptor cDNA (65%), which was previously cloned from rat uterus (Koike, S., Sakai, M., & Muramatsu, M. (1987) *Nucleic Acids Res* 15, 2499-2513). The amino acid residues predicted by clone 29 DNA sequences suggested that this DNA fragment encoded part of the DNA-binding domain, hinge region and the beginning of the ligand binding domain of a novel member of the nuclear receptor family. Two PCR primers (Figure 1) were used to generate a probe of 204 bp consisting of the hinge region of the novel receptor, which was used to screen a rat prostate cDNA library (Clontech gt10) under stringent conditions resulting in four strongly positive clones with a size of 0.9 kb, 1.8kb, 2.5kb and 5-6kb respectively. The clone of 2.5kb was sequenced and Figure 1 shows the nucleotide sequence determined in the core facility (CyberGene AB) by cycle sequencing using fluorescent terminators (Applied Biosystems) on both strands, with a series of internal primers and deduced amino acid sequence of clone 29. Two in frame ATG codons are located at nucleotide 424 and nucleotide 448, preceding by an in-frame stop codon at nucleotide 319, which suggests that they are possible start codons. The open reading frame encodes a protein of 485 amino acid residues (counted from the first methionine) with a calculated molecular weight of 54.2 kDa. Analysis of the proteins synthesized by *in-vitro* translation from the clone 29 cRNA in rabbit reticulocyte lysate revealed a doublet protein band migrating at approximately 57 kDa on SDS-PAGE gels (data not shown), confirming the open reading frame. The doublet protein band is probably caused by the use of both ATG codons for initiation of protein synthesis. The amino acid sequence of clone 29 protein shows the characteristic zinc module DNA-binding domain, hinge region and a putative ligand binding domain, which are the characteristic features of members of the nuclear receptor family (Tsai, M.-J., & O'Malley, B.W (1994) *Ann. Rev. Biochem.* 63, 451-486; Hård, T., & Gustafsson, J.-Å (1993) *Acc. Chem. Res.* 26, 644-650; Laudet, V.; Hänni, C., Coli, J., Catzeflis, F., & Stehelin, D (1992) *EMBL J.* 11, 1003- 1012).

[0014] Protein sequence comparison with several representative members of the nuclear receptor family (Figure 2) showed the clone 29 protein is most related to the rat estrogen receptor (ER $\alpha$ ), cloned from uterus (Koike, S., Sakai, M., & Muramatsu, M. (1987) *Nucleic. Acids Res.* 15, 2499-2513), with 95% identity in the DNA-binding domain (amino acid residues 103-167) (Griffiths, K., Davies, P., Eaton, C.I., Harper, M.E., Turkes, A., & Peeling, W.B. (1991) in *Endocrine Dependent Tumours*, eds. Voigt, K.-D. & Knabbe, C. (Raven Press), pp. 83-125). A number of functional characteristics have been identified within the DNA-binding domain of nuclear receptors (Hard, T., & Gustafsson, J.-Å, (1993) *Acc. Chem. Res.* 26, 644-650 and Zilliacus, J., Carlstedt-Duke, J., Gustafsson, J.-Å, & Wright, A.P.H. (1994) *Proc. Natl. Acad. Sci. USA* 91, 4175-4179). The so-called P-box specifies nucleotide sequence recognition of the core half-site within the response element, while the D-box mediates dimerization between receptor monomers. The clone 29 protein P-box and D-box sequences of EGCKA and PATNQ, respectively, are identical to the corresponding boxes in ER $\alpha$  (Hård, T., & Gustafsson, J.-Å, (1993) *Acc. Chem. Res.* 26, 644-650 and Koike, S., Sakai, M., & Muramatsu, M. (1987) *Nucleic Acids Res.* 15, 2499-2513), thus predicting that clone 29 protein binds to ERE sequences.

[0015] The putative ligand binding domain (LBD) of clone 29 protein (amino acid residues 259-457) shows closest homology to the LBD of the rat ER $\alpha$  (Figure 2), while the homology with the human ERR1 and ERR2 proteins (Giguere, V., Yang, N., Segui, P., & Evans R.M. (1988) *Nature* 331, 91-94) is considerably less. With the human, mouse and xenopus estrogen receptors the homology in the LBD is also around 55%, while the homology with the LBD of other steroid receptors is not significant (Figure 2). Cysteine residue 530 in human ER $\alpha$  has been identified as the covalent attachment site of an estrogenic affinity label (Harkow, K.W., Smith D.N., Katzenellenbogen, J.A., Greene, G.L., & Katzenellenbogen, B.S. (1989) *J. Biol. Chem.* 264, 17476- 17485). Interestingly, clone 29 protein (Cys-436) as well as the mouse, rat and xenopus ER $\alpha$ s have a cysteine residue at the corresponding position. Also, two other amino acid residues described to be close to or part of the ligand-binding pocket of the human ER $\alpha$ -LBD (Asp 426 and Gly 521) are conserved in the LBD of clone 29 protein (Asp 333 and Gly 427) and in the LBD of ER $\alpha$ s from various species (20.21). The ligand-dependent transactivation function TAF-2 identified in ER $\alpha$  (Danielian, P.S., White, R., Lees, J.A., & Parker, M.G. (1992) *EMBO J.* 11, 1025-1033), which is believed to be involved in contacting other transcription factors and thereby influencing activation of target genes, is almost completely conserved in clone 29 protein (amino acid residues 441-457). Steroid hormone receptors are phosphoproteins (Kuiper, G., & Brinkmann, A. O. (1994) *Mol. Cell. Endocrinol.* 100, 103-107), and several phosphorylation sites identified in the N-terminal domain and LBD of ER $\alpha$  (Arnold, S.F., Obourn, J.D., Jaffe, H., & Notides, A.C. (1995) *Mol. Endocrinol.* 9, 24-33 and Le Goff, P., Montano, M.M., Schodin, D.J., & Katzenellenbogen, B.S (1994) *J. Biol. Chem.* 269, 4458-4466) are conserved in clone 29 protein (Ser 30 and 42, Tyr 443). Clone 29 protein consists of 485 amino acid residues while ER $\alpha$ s from



human, mouse and rat consist of 590-600 amino acid residues. The main difference is a much shorter N-terminal domain in clone 29 protein i.e 103 amino acid residues as compared to 185-190 amino acid residues in the other receptor proteins. Also the non-conserved so-called F-domain at the C-terminal end of ER $\alpha$ s is 15 amino acid residues shorter in clone 29 protein. The cDNA insert of a positive clone of 2.6 kb was subcloned into the EcoR1 site of pBluescript (trademark) (Stratagene). The complete DNA sequence of clone 29 was determined (CyberGene AB) by cycle sequencing using fluorescent terminators (Applied Biosystems) on both strands, with a series of internal primers. [0016] Figs 2C and 2D respectively compare the ligand and DNA binding domain of Er $\beta$  compared to rat, mouse and human Er $\alpha$ 's.

## 2. Saturation ligand binding analysis and ligand competition studies:

[0017] Clone 29 cDNA was subcloned in pBluescript downstream of the T7 promoter to give p29-T7. Clone 29 protein was synthesized *in vitro* using the TnT-coupled reticulocyte lysate system (Promega). Translation reaction mixtures were diluted five times with TEDGMO buffer (40 mM Tris/HCl, pH 7.4, 1 mM EDTA, 10% (v/v) glycerol, 10 mM Na<sub>2</sub>MoO<sub>4</sub>, 10 mM DTT) and 0.1 ml aliquots were incubated for 16 h at 8° C with 0.3- 6.2 nM [2,4,6,7-<sup>3</sup>H]-17 $\beta$ -estradiol (NEN-Dupont; specific radioactivity 85 Ci/mmol) in the presence or absence of a 200-fold excess of unlabelled E2.

[0018] Fig. 5A illustrates the results of a saturation ligand analysis of clone 29 protein. Reticulocyte lysate containing clone 29 protein was incubated with 6 concentrations of [<sup>3</sup>H]E2 between 0.3 and 6.0 nM. Parallel tubes contained an additional 200 fold of non-radioactive E2. Bound and free ligand were separated with a dextran-coated charcoal assay. The K<sub>d</sub> (0.6 nM) was calculated from the slope of the line in the Scatchard plot shown ( $r = 0.93$ ), and the number of binding sites was extrapolated from the intercept on the abscissa ( $B_{max} = 1400$  fmol/ml undiluted translation mixture).

[0019] For ligand competition studies diluted reticulocyte lysate was incubated with 5 nM [2,4,6,7-<sup>3</sup>H]-17 $\beta$ -estradiol in the presence of either 0, 5, 50, 500 or 5,000 nM of non-radioactive E2, estrone, estriol, testosterone, progesterone, corticosterone, 5 $\alpha$ -androstane-3 $\beta$ ,17 $\beta$ -diol, 5 $\alpha$ -androstane-3 $\alpha$ ,17 $\beta$ -diol and diethylstilbestrol (DES) for 16 h at 8° C. Bound and unbound steroids were separated with a dextran-coated charcoal assay (Ekman, P., Barrack, E.R., Greene, G.L., Jensen, E.V., & Walsh, P.C (1983) J. Clin. Endocrinol Metab. 57, 166-176).

[0020] Fig. 5B illustrates the specificity of ligand binding by clone 29 protein. Reticulocyte lysate containing clone 29 protein was equilibrated for 16 h with 5 nM [<sup>3</sup>H]E2 and the indicated fold excess of competitors. Data represent [<sup>3</sup>H]E2 bound in the presence of unlabelled E2, testosterone (T), progesterone (prog), corticosterone (cortico), estrone (E1), diethylstilbestrol (DES), 5 $\alpha$ -androstane-3 $\alpha$ , 17 $\beta$ -diol (3 $\alpha$ -AD), 5 $\alpha$ -androstane-3 $\beta$ ,17 $\beta$ -diol (3 $\beta$ -AD) and estriol (E3). [<sup>3</sup>H]E2 binding in the absence of competitor was set at 100%.

## 3. In-situ hybridisation:

[0021] *In-situ* hybridisation was carried out as previously described (Dagerlind Å., Friberg, K., Bean, A.J., & Hökfelt, T (1992) *Histochemistry* 98, 39-49). Briefly, two oligonucleotide probes directed against nucleotides 994-1041 and 1981-2031 were each labelled at the 3'-end with <sup>32</sup>P-dATP using terminal deoxynucleotidyltransferase (Amersham, UK). Adult male and female Sprague-Dawley rats (age 2 to 3 months  $n=10$ ) were used for this study. The rats were decapitated and the tissues were rapidly excised and frozen on dry ice. The tissues were sectioned in a Microm HM500 cryostat at 14  $\mu$ m and thawed onto Probe-On glass slides (Fisher Scientific, PA, USA). The slides were stored at -20° C until used. The slides were incubated in humidified boxes at 42° C for 18 h with  $1 \times 10^6$  cpm of the probe in a hybridization solution containing 50% formamide, 4 x SSC (1 x SSC = 0.15 M NaCl, 0.015 M sodium citrate), 1 x Denhardt (0.02 % BSA, 0.02 % Ficoll, 0.02 % PVP), 1 % sarkosyl, 0.02 M sodium phosphate (pH 7.), 10% dextran sulphate, 500  $\mu$ g/ml salmon sperm DNA and 200 mM DTT. Slides were subsequently rinsed in 1 x SSC at 55° C for 60 min with four changes of SSC and finally in 1 x SSC starting at 55° C and slowly cooled to room temperature, transferred through distilled water and briefly dehydrated in 50% and 95% ethanol for 30 sec each, air-dried, and covered with Amersham  $\beta$ -man autoradiography film for 15 to 30 days. Alternatively the slides were dipped in Kodak NTB2 nuclear track emulsion (diluted 1:1 with distilled water) and exposed for 30 to 60 days at 4° C. Finally, the sections were stained with cresyl violet.

[0022] Clear expression of clone 29 was observed in the reproductive tract of both male and female rats, while in all other rat tissues the expression was very low or below the level of detection with *in-situ* hybridisation (not shown). In male reproductive organs high expression was seen in the prostate gland (Figure 3), while very low expression was observed in testis, epididymis and vesicula seminalis (not shown). In dipped sections, expression was clearly visible in prostate epithelial cells (secreting alveoli) while the expression in smooth muscle cells and fibroblasts in the stroma was low (Figure 3). In female reproductive organs expression was seen in the ovary (Figure 4), while uterus and vagina were negative (not shown). In dipped sections high expression was seen in the granulosa cell layer of primary, secondary and mature follicles (Figure 4), whereas primordial follicles, oocytes and corpora lutea appeared completely negative. Low expression was seen in the interstitial cells of the ovary. Both anti-sense oligonucleotide probes used produced similar results. Addition of a 100 fold excess of the respective unlabelled oligonucleotide probes during the



hybridisation reactions abolished all signals.

#### 4. Transactivation analysis in CHO-cells:

[0023] The expression vector pCMV29 was constructed by inserting the 2.6 kb clone 29 fragment in the EcoRI site of the expression vector pCMV5 (Andersson, S., Davis, D.L., Dahlbäck, H., Jörnvall, H., & Russell, D.W. (1989) *J. Biol. Chem.* **264**, 8222-8229). The pERE-ALP reporter construct contains a secreted form of the placental alkaline phosphatase gene (Berger, J., Hauber, J., Hauber, R., Geiger, R., & Cullen, B.R. (1988) *Gene* **66**, 1-10) and the MMTV-LTR in which the glucocorticoid response elements were replaced by the vitellogenin promoter estrogen response element (ERE).

[0024] CHO-K1 cells were seeded in 12-well plates at approximately  $1.7 \times 10^5$  cells per well in phenol-red free Ham F12 medium with 5% FCS (dextran-coated charcoal treated) and 2 mM L-glutamine. After 24 h the cells were transfected with 250 ng pERE-ALP vector and 50 ng pCMV29 using lipofectamine (Gibco) according to the manufacturer's instructions. After five hours of incubation the cells were washed and refed with 0.5 ml phenol-red free Coon's F-12 medium containing 5% serum substitute (SRC 3000, Tissue Culture Services Ltd., Botolph Claydon, Buckingham, UK) 2 mM L-glutamine and 50 µg/ml gentamicin plus hormones as indicated. After 48 h the medium was assayed for alkaline phosphatase (ALP) activity by a chemiluminescence assay. A 10 µl aliquot of the cell culture medium was mixed with 200 µl assay buffer (10 mM diethanolamine pH 10.0 1 mM  $MgCl_2$  and 0.5 mM CSPD (Tropix Inc. Boston, USA)) and incubated for 20 min at 37°C before measurement in a microplate luminometer (Luminoskan; Labsystems, Finland) with integral measurement for 1 second. The ALP activity of ERE-reporter alone was set at 1.

#### 5. Ligand binding characteristics and transactivation function of clone 29 protein:

[0025] On the basis of the described high homology between clone 29 protein and rat ERα in the DBD and LBD it was hypothesized that clone 29 protein might encode a novel ER. Furthermore, biological effects of estrogens on rat prostate and ovary, which show high expression of clone 29 RNA, are well known (Griffiths, K., Davies, P., Eaton, C. I., Harper, M.E., Turkes, A., & Peeling W. B. (1991) in *Endocrine Dependent Tumours*, eds Voigt, K-D. & Khabbe, C. (Raven Press), pp 83-125; Richards, J.S (1994) *Endocrine Rev.* **15**, 725-745; and Habenicht, U-F., Tunn, U.W., Senge, Th., Schroder, R.H., Schweikert, H.U., Bartsch, G., & El Etreby, M.F. (1993) *J. Steroid Biochem. Molec. Biol.* **44**, 557-563). In order to analyze the steroid binding properties of clone 29 protein synthesized *in vitro*, the reticulocyte lysate was incubated at 8°C for 16 h with increasing concentrations (0.3-6.0 nM) of [ $^3H$ ]E2 in the presence or absence of a 200 fold molar excess of unlabelled E2. Linear transformation of saturation data revealed a single population of binding sites for E2 with a  $K_d$  (dissociation constant) of 0.6 nM (Figure 5A and C). Steroid binding specificity was measured by incubating reticulocyte lysate with 5 nM [ $^3H$ ]E2 in the presence of 0.5, 50, 500 and 5,000 nM unlabelled competitors. Competition curves generated are indicative of an estrogen receptor in that only estrogens competed efficiently with [ $^3H$ ]E2 for binding (Figure 5B). Fifty percent inhibition of specific binding occurred by 0.6 fold excess of unlabelled E2; diethylstilbestrol, estriol, estrone and 5α-androstane-3β,17β-diol were 5, 15, 50 and 150 times, respectively, less effective as competitors. Neither testosterone, progesterone, corticosterone nor 5α-androstane-3α,17β-diol were efficient competitors, even at the highest concentrations used (1000 fold excess). The dissociation constant and the steroid binding specificities measured are in good agreement with data previously reported for ERs in rat and human prostate, rat granulosa cells, rat antral follicles and whole rat ovarian tissue (Ekman, P., Barrack, E.R., Greene, G.L., Jensen, E.V., & Walsh, P.C (1983) *J. Clin. Endocrinol. Metab.* **57**, 166-176; van Beurden-Lamers, W.M.O., Brinkmann, A-O., Mulder, E., & van der Molen, H. (1974) *Biochem. J.* **140**, 495-502; Kudolo, G.B., Elder, M.G., & Myatt, L. (1984) *J. Endocrinol.* **102**, 83-91; and Kawashima, M., & Greenwald, G.S. (1993) *Biology of Reprod.* **48** 172-179).

[0026] When clone 29 protein was labelled with a saturating dose of [ $^3H$ ]E2 and analyzed on sucrose density gradients, a single peak of specifically bound radioactivity was observed. The sedimentation coefficient of this complex was about 7S, and it shifted to 4S in the presence of 0.4 M NaCl (not shown). To investigate the transcriptional regulatory properties of clone 29 protein, we performed co-transfection experiments in which CHO cells were transfected with a clone 29 protein expression vector and/or an estrogen-responsive reporter gene construct. Cells were incubated in the absence of E2 (clone 29) or in the presence of 100 nM E2 (Clone 29 + E2) or in the presence of 100 nM E2 and 12 µM Tamoxifen (Clone 29 + E2/Tam). In the absence of exogenously added E2 clone 29 protein showed considerable transcriptional activity which could be further increased by the addition of 100 nM E2 (Figure 6). Simultaneous addition of a 10 fold excess of the antiestrogen Tamoxifen partially suppressed the E2 stimulated activity (Figure 6). The constitutive transcriptional activity of clone 29 protein could be suppressed by the anti-estrogen ICI-1624384 (not shown). It has been shown previously that the wild-type mouse and human ERs are constitutive activators of transcription, and that the transcriptional activity can be stimulated further by the addition of E2 (Txukerman, M., Xiao-Kun Zhang, Her-mann, T., Wills, K. N., Graupner, G., & Phal, M. (1990) *New Biologist* **2**, 613-620 and Lees, J.A., Fawell, S.E., & Parker, M.G. (1989) *Nucl. Acids Res.* **17**, 5477-5488). To obtain more insight into what concentrations of E2 effect clone 29



protein transcriptional activity, transient transfection experiments were carried out in the presence of increasing concentrations of E2. CHO-cells were transiently transfected with the ERE-reporter plasmid and the clone 29 protein expression plasmid. Cells were incubated with increasing concentrations of E2 (0.1 - 1000 nM), estrone (E1, 1000 nM), 5 $\alpha$ -androstane-3 $\beta$ ,17 $\beta$ -diol (3 $\beta$ -AD, 1000 nM) or no ligand added. Alkaline phosphatase activity (ALP) was measured as described and the activity in the absence of ligand (control) was set at 1. The figure shows relative ALP-activities ( $\pm$ SD) from three independent experiments. Clone 29 protein began to respond at 0.1 nM E2 and maximal stimulation was observed between 1 nM and 10 nM E2 (Figure 7). The maximal stimulation factor was  $2.6 \pm 0.5$  fold (mean  $\pm$  SD, n = 9) as compared to incubation in the absence of E2. Apart from E2 also estrone and 5 $\alpha$ -androstane-3 $\beta$ ,17 $\beta$ -diol could stimulate transcriptional activity, albeit at higher concentrations (Figure 7). Dexamethasone, testosterone, progesterone, 5 $\alpha$ -androstane-3 $\alpha$ ,17 $\beta$ -diol, thyroid hormone and all-*trans*-retinoic acid could not stimulate transcriptional activity of clone 29 protein, even at the highest concentration (1000 nM) tested (not shown). The results of the co-transfection experiments are in agreement with the ligand binding and specificity data of clone 29 protein presented in Figure 5. In control experiments, wild-type human ER $\alpha$  also showed transcriptional activity in the absence of E2, which could be increased by the addition of E2 (not shown).

#### 6. Detection of rat ER expression by RT-PCR

[0027] The tissue specificity of expression of rat ER $\beta$  and ER $\alpha$  was determined using reverse transcriptase polymerase chain reaction (RT-PCR). The results of the experiment are shown in Fig. 8.

#### B. Isolation of human Er $\beta$

[0028] 1. A human version of Er $\beta$  (hER $\beta$ ) has also been cloned from human ovary. The tissue specificity of hER $\beta$  expression in a variety of cells was also determined using the RT-PCR technique. The results are shown in Fig. 9. It will be noticed that there is a very high level of mRNA of hER $\beta$  in human umbilical vein endothelial cells (HUVEC) but no detection of hER $\alpha$  in the same cells. In addition, it will be seen that in human osteosarcoma cell line (HOS-D4), hER $\beta$  is expressed in greater quantities compared to hER $\alpha$ .

[0029] 1. A human version of ER $\beta$  (hER $\beta$ ) has also been cloned. The tissue specificity of hER $\beta$  expression in a variety of cells was also determined using the RT-PCR technique. The results are shown in Fig. 9. It will be noticed that there is a very high level of mRNA of hER $\beta$  in human umbilical vein endothelial cells (HUVEC) but no detection of hER $\alpha$  in the same cells. In addition, it will be seen that in human osteosarcoma cell line (HOS-D4), hER $\beta$  is expressed in greater quantities compared to hER $\alpha$ .

[0030] The partial DNA sequence of hER $\beta$  is shown in Fig. 10 and a derived amino acid sequence is shown in Fig. 11.

#### Cloning of human Er $\beta$ from testis

[0031] A commercially available cDNA from human testis (Clontech, article no. HL1161x) was screened, using a fragment containing the ligand-binding domain of the rat Er $\beta$  cDNA as probe. Approximately  $10^6$  recombinants were screened, resulting in one positive clone. Upon sequencing of this clone, it was seen that the insert was 1156 bp (Figure 13A and 13B). This corresponds to most of the translated region of a receptor with an overall homology of 90.0% to rat Er $\beta$ , therefore deduced to represent the human form of Er $\beta$ .

[0032] The cloned hER $\beta$ , however, lacks approximately 47 amino acids at the N-terminal end and 61 amino acids at the C-terminal end (as compared to the rat sequence). Further screening of the same library was unsuccessful. PCR technology was therefore used to obtain the remaining parts. For oligonucleotides were synthesised; two degenerate oligonucleotides containing all possible codons for the amino acids adjacent to the initiation methionine and the stop codon, respectively, of the rat Er $\beta$ , and two specific oligonucleotides containing the sequence of the clone isolated from the human testis library and situated approximately 100 bp from respective end of this clone. PCR with the N-terminal and C-terminal pair of oligos yielded specific bands, that were subcloned and sequenced. The parts of these new clones that overlap the original cDNA clone are identical to this. It was thus possible to construct peptide and DNA sequences corresponding to the whole open reading frame (Fig. 13A and 13B).

[0033] When comparing the human Er $\beta$  to rat Er $\beta$ , this receptor is 79.6% identical in the N-terminal domain, 98.5% in the DNA-binding domain, 85.6% in the hinge and 91.6% in the ligand-binding and F-domains. These numbers match very well those found when comparing the rat and human forms of Era.

[0034] Studies of the expression of human Er $\beta$  using Northern blot show expression in testis and in ovaries. The expression in prostate, however, appears lower than found in the rat.

[0035] The human Er $\beta$  gene has been mapped to chromosome 14 using PCR and to region 14q22-23 using the FISH technique, whereas the human Er $\beta$  gene has been mapped to chromosome 6q25.



## 2. Comparison of ligand binding affinity of hER $\alpha$ and rER $\beta$

[0036] The ligand affinity of the two estrogen receptors, human ER $\alpha$  (ovary) (hER $\alpha$ ) and rat ER $\beta$  (rER $\beta$ ) was tested in binding saturation experiments and in binding competition experiments.

[0037] cDNA of the receptor subtypes hER $\alpha$  and rER $\beta$  were *in vitro* translated in rabbit reticulocyte lysate in presence of non-radioactive amino acids according to the instructions supplied by the manufacturer (Promega).

[0038] The radioactive ligand used in all experiments was 16 $\alpha$ -[<sup>125</sup>I]-17 $\beta$ -estradiol ([<sup>125</sup>I]-E2) (NEX- 144, New England Nuclear). The method for the binding experiments was previously described in: Salomonsson M, Carlsson B, Haggblad J. J. *Steroid Biochem. Molec. Biol.* Vol. 50, No. 5/6 pp. 313-18, 1994. In brief, estrogen receptors are incubated with [<sup>125</sup>I]-E2 to equilibrium (16-18 h at +4°C). The incubation was stopped by separation of protein-bound [<sup>125</sup>I]-E2 from free [<sup>125</sup>I]-E2 on Sephadex G25 columns. The radioactivity of the eluate is measured in a gamma-counter.

[0039] In the competition experiments, non-radioactive ligands were diluted in DMSO, mixed with [<sup>125</sup>I]-E2 (approximately 100-200 pM), aliquoted in parallel, and finally hER $\alpha$  or rER $\beta$  was added. The final concentration of DMSO in the binding buffer was 2%.

[0040] The buffer used in the experiments was of the following composition: Hepes (pH=7.5) 20 mM, KCl 150 mM, EDTA 1 mM, glycerol (8.7%), monothioglycerol 6 mM, Na<sub>2</sub>MO<sub>4</sub> 10mM.

## 3. Equilibrium binding saturation experiments (K<sub>d</sub>-determinations)

[0041] A range of concentrations of [<sup>125</sup>I]-E2 were mixed with the ER:s and incubated as described above, free [<sup>125</sup>I]-E2 was determined by subtracting bound [<sup>125</sup>I]-E2 from added [<sup>125</sup>I]-E2. Binding data was analysed by Hill-plots and by Scatchard plots (Figure 11). The equilibrium binding results are shown in Table 1. The apparent K<sub>d</sub>-values for [<sup>125</sup>I]-E2 differed between the two ER:s with approximately a factor of four: K<sub>d</sub>(hER $\alpha$ ):K<sub>d</sub>(rER $\beta$ ) = 1:4.

Table 1.

Equilibrium dissociation constants for [ <sup>125</sup> I]-E2 to the two subtypes.		
Receptor subtype	K <sub>d</sub> (Hill-plot)	K <sub>d</sub> (Scatchard-plot)
hER $\alpha$	0.06 nM	0.09 nM
rER $\beta$	0.24 nM	0.42 nM

## 4. Competition experiments (IC<sub>50</sub> determinations)

[0042] The experiments were performed as described above. IC<sub>50</sub> values were obtained by applying a four parameter logistic analysis;  $b = ((b_{max} - b_{min}) / (1 + (1/IC_{50})^S)) + b_{min}$ , where I is the added concentration of binding inhibitor, IC<sub>50</sub> is the concentration of inhibitor at half maximal binding and S is a slope factor. The free concentration of [<sup>125</sup>I]-E2 was determined by sampling an aliquot from the wells at the end of the incubation and then subtract bound radioactivity from sampled total radioactivity.

[0043] Since the equilibrium binding experiments (above) showed that the K<sub>d</sub>-values for [<sup>125</sup>I]-E2 differed between the two ER:s, K<sub>i</sub>-values (from the Cheng-Prusoff equation:  $K_i = IC_{50} / (1 + L/K_d)$  where L is free ([<sup>125</sup>I]-E2)) were calculated for the compounds investigated. Two approaches for calculating RBA (Relative Binding Affinity) were used. The RBA values were derived using either the IC<sub>50</sub> values or the K<sub>i</sub> values. In both approaches, the value for the compound 16 $\alpha$ -bromo-estradiol was selected as the reference value (100%). Both approaches gave similar results. The results are summarized in Figure 12. In these Figures "4-OH-Tam" = 4-hydroxy-tamoxifen; "DES" = diethylstilbestrol; "Hexestr" = hexestrol; "ICI-164384" = ICI plc compound no. 164382; "17 $\beta$ -E2" = 17 $\beta$ -estradiol; "16 $\alpha$ -B-E2" = 16 $\alpha$ -bromo-estradiol; "Ralox" = Raloxifen; and "17 $\alpha$ -E2" 17 $\alpha$  diol.

[0044] The results show that ER $\alpha$  and ER $\beta$  have significant different ligand binding affinities - the apparent K<sub>d</sub>-values for [<sup>125</sup>I]-E2 differed between the two ER's by a factor of about 4 (K<sub>d</sub>(hER $\alpha$ ): K<sub>d</sub>(rER $\beta$ )  $\approx$  1:4). Some compounds investigated showed significant differences in the competition for binding of [<sup>125</sup>I]-E2 to the ER's. Certain compounds were found to be more potent inhibitors of [<sup>125</sup>I]-E2 binding to hER $\alpha$  as compared to rER $\beta$  whereas others were found to be more potent inhibitors of [<sup>125</sup>I]-E2 binding to rER $\beta$  than to hER $\alpha$ .

## Claims

1. An isolated estrogen receptor (called ER $\beta$ ) having the amino acid sequence of Fig. 1.



EP 0 792 292 B1

2. An isolated estrogen receptor (called ER $\beta$ ) having an amino acid sequence which is more than 95% identical to the amino acid sequence of Fig. 1.
3. An isolated estrogen receptor (called ER $\beta$ ) having the amino acid sequence of Fig. 13A.
- 5 4. An isolated estrogen receptor (called ER $\beta$ ) having an amino acid sequence which is more than 89% identical to the amino acid sequence of Fig. 13A, and which is expressed in greater quantities in HUVEC and HOS D4 cells than ER $\alpha$ .
- 10 5. An isolated estrogen receptor (called ER $\beta$ ) having the amino acid sequence of Fig. 14A.
6. An isolated estrogen receptor (called ER $\beta$ ) having an amino acid sequence which is more than 95% identical to the amino acid sequence of Fig. 14A.
- 15 7. An estrogen receptor according to any one of claims 1 to 6 which is derived from mammalian cells.
8. An estrogen receptor according to claim 7 which is derived from rat or human cells.
9. An isolated DNA sequence encoding an estrogen receptor according to any preceding claim.
- 20 10. An isolated DNA sequence according to claim 9 in which the DNA sequence is one of those given in Figs. 1, 13B or 14B.
- 25 11. An isolated nucleic acid sequence which is capable of hybridizing under stringent conditions to an oligonucleotide probe comprising nucleotides 994-1041 of Fig. 1, wherein said nucleic acid sequence codes for an estrogen receptor according to any one of claims 1 to 8.
- 30 12. An isolated nucleic acid sequence which is capable of hybridizing under stringent conditions to an oligonucleotide probe comprising nucleotides 1981-2031 of Fig. 1, wherein said nucleic acid sequence codes for an estrogen receptor according to any one of claims 1 to 8.
- 35 13. The use of an estrogen receptor according to any one of claims 1 to 8, or a DNA sequence according to claim 9 or 10, or a nucleic acid sequence according to claim 11 or 12 in an assay to determine molecules which bind an estrogen receptor according to any one of claims 1 to 8.
- 40 14. The use of an estrogen receptor according to any one of claims 1 to 8, or a DNA sequence according to claim 9 or 10, or a nucleic acid sequence according to claim 11 or 12 in an assay to determine molecules for use in the treatment of ER $\beta$  specific diseases or conditions or diseases or conditions related to an estrogen receptor according to any one of claims 1 to 8.
- 45 15. The use of an estrogen receptor according to any one of claims 1 to 8, or a DNA sequence according to claim 9 or 10, or a nucleic acid sequence according to claim 11 or 12 in an assay to determine molecules for use in the treatment of prostate or ovarian cancer, benign prostatic hyperplasia, diseases of the central nervous system, osteoporosis, or cardiovascular disease.
- 50 16. A drug design method comprising comparing binding of a test compound to ER $\alpha$  and to an estrogen receptor according to any one of claims 1 to 8.
17. The use of an estrogen receptor according to any one of claims 1 to 8 in the testing of the possible estrogenic or other hormonal effects of a substance.

Patentansprüche

- 55 1. Isolierter Östrogenrezeptor (benannt ER $\beta$ ) mit der Aminosäuresequenz von Fig. 1.
2. Isolierter Östrogenrezeptor (benannt ER $\beta$ ) mit einer Aminosäuresequenz, die zu mehr als 95 % identisch ist mit der Aminosäuresequenz von Fig. 1.



3. Isolierter Östrogenrezeptor (benannt ER $\beta$ ) mit der Aminosäuresequenz von Fig. 13A.
4. Isolierter Östrogenrezeptor (benannt ER $\beta$ ) mit einer Aminosäuresequenz, die zu mehr als 89 % identisch ist mit der Aminosäuresequenz von Fig. 13A und die in HUVEC- und HOS D4-Zellen in größeren Mengen als ER $\alpha$  exprimiert wird.
5. Isolierter Östrogenrezeptor (benannt ER $\beta$ ) mit der Aminosäuresequenz von Fig. 14A.
6. Isolierter Östrogenrezeptor (benannt ER $\beta$ ) mit einer Aminosäuresequenz, die zu mehr als 95 % identisch ist mit der Aminosäuresequenz von Fig. 14A.
7. Östrogenrezeptor nach einem der Ansprüche 1 bis 6, der von Säugerzellen stammt.
8. Östrogenrezeptor nach Anspruch 7, der von Ratten- oder menschlichen Zellen stammt.
9. Isolierte DNA-Sequenz, die einen Östrogenrezeptor nach einem der vorangehenden Ansprüche codiert.
10. Isolierte DNA-Sequenz nach Anspruch 9, in der die DNA-Sequenz eine der Sequenzen ist, die in den Figuren 1, 13B oder 14B gezeigt ist.
11. Isolierte Nucleinsäuresequenz, die unter stringenten Bedingungen mit einer Oligonucleotidsonde hybridisieren kann, die die Nucleotide 994-1041 von Fig. 1 umfaßt, wobei die Nucleinsäuresequenz einen Östrogenrezeptor nach einem der Ansprüche 1 bis 8 codiert.
12. Isolierte Nucleinsäuresequenz, die unter stringenten Bedingungen mit einer Oligonucleotidsonde hybridisieren kann, die die Nucleotide 1981-2031 von Fig. 1 umfaßt, wobei die Nucleinsäuresequenz einen Östrogenrezeptor nach einem der Ansprüche 1 bis 8 codiert.
13. Verwendung eines Östrogenrezeptors nach einem der Ansprüche 1 bis 8 oder einer DNA-Sequenz nach Anspruch 9 oder 10 oder einer Nucleinsäuresequenz nach Anspruch 11 oder 12 in einem Test zur Bestimmung von Molekülen, die an einen Östrogenrezeptor nach einem der Ansprüche 1 bis 8 binden.
14. Verwendung eines Östrogenrezeptors nach einem der Ansprüche 1 bis 8 oder einer DNA-Sequenz nach Anspruch 9 oder 10 oder einer Nucleinsäuresequenz nach Anspruch 11 oder 12 in einem Test zur Bestimmung von Molekülen zur Verwendung in der Behandlung von ER $\beta$ -spezifischen Krankheiten oder Zuständen oder Krankheiten oder Zuständen, die mit einem Östrogenrezeptor nach einem der Ansprüche 1 bis 8 im Zusammenhang stehen.
15. Verwendung eines Östrogenrezeptors nach einem der Ansprüche 1 bis 8 oder einer DNA-Sequenz nach Anspruch 9 oder 10 oder einer Nucleinsäuresequenz nach Anspruch 11 oder 12 in einem Test zur Bestimmung von Molekülen zur Verwendung in der Behandlung von Prostata- oder Eierstockkrebs, gutartiger Prostatahyperplasie, Krankheiten des zentralen Nervensystems, Osteoporose oder Herz-Kreislauf-Erkrankungen.
16. Verfahren zur Konstruktion eines Wirkstoffes, umfassend das Vergleichen der Bindung einer Testverbindung an ER $\alpha$  und an einen Östrogenrezeptor nach einem der Ansprüche 1 bis 8.
17. Verwendung eines Östrogenrezeptors nach einem der Ansprüche 1 bis 8 zum Testen der möglichen Östrogen- oder anderer hormonaler Wirkungen eines Stoffes.

#### Revendications

1. Récepteur d'estrogènes isolé (appelé ER $\beta$ ) ayant la séquence d'acides aminés de la figure 1.
2. Récepteur d'estrogènes isolé (appelé ER $\beta$ ) ayant une séquence d'acides aminés identique à plus de 95 % à la séquence d'acides aminés de la figure 1.
3. Récepteur d'estrogènes isolé (appelé ER $\beta$ ) ayant la séquence d'acides aminés de la figure 13A.



4. Récepteur d'estrogènes isolé (appelé ER $\beta$ ) ayant une séquence d'acides aminés identique à plus de 89 % à la séquence d'acides aminés de la figure 13A, et qui est exprimé en de plus grandes quantités dans les cellules HUVEC (cellules endothéliales de la veine ombilicale humaine) et HOS-D4 (lignée cellulaire d'ostéosarcome humain) que l'ER $\alpha$ .
5. Récepteur d'estrogènes isolé (appelé ER $\beta$ ) ayant la séquence d'acides aminés de la figure 14A.
6. Récepteur d'estrogènes isolé (appelé ER $\beta$ ) ayant une séquence d'acides aminés identique à plus de 95 % à la séquence d'acides aminés de la figure 14A.
7. Récepteur d'estrogènes selon l'une quelconque des revendications 1 à 6, qui est dérivé de cellules de mammifères.
8. Récepteur d'estrogènes selon la revendication 7, qui est dérivé de cellules de rat ou de cellules humaines.
9. Séquence d'ADN isolée codant pour un récepteur d'estrogènes selon l'une quelconque des revendications précédentes.
10. Séquence d'ADN isolée selon la revendication 9, dans laquelle la séquence d'ADN est l'une de celles données dans les figures 1, 13B ou 14B.
11. Séquence d'acides nucléiques isolée, qui est capable de s'hybrider dans des conditions stringentes à une sonde oligonucléotidique comprenant les nucléotides 994-1041 de la figure 1, ladite séquence d'acides nucléiques codant pour un récepteur d'estrogènes selon l'une quelconque des revendications 1 à 8.
12. Séquence d'acides nucléiques isolée, qui est capable de s'hybrider dans des conditions stringentes à une sonde oligonucléotidique comprenant les nucléotides 1981-2031 de la figure 1, ladite séquence d'acides nucléiques codant pour un récepteur d'estrogènes selon l'une quelconque des revendications 1 à 8.
13. Utilisation d'un récepteur d'estrogènes selon l'une quelconque des revendications 1 à 8, ou d'une séquence d'ADN selon la revendication 9 ou 10, ou d'une séquence d'acides nucléiques selon la revendication 11 ou 12, dans une analyse destinée à la détermination de molécules qui se lient à un récepteur d'estrogènes selon l'une quelconque des revendications 1 à 8.
14. Utilisation d'un récepteur d'estrogènes selon l'une quelconque des revendications 1 à 8, ou d'une séquence d'ADN selon la revendication 9 ou 10, ou d'une séquence d'acides nucléiques selon la revendication 11 ou 12, dans une analyse destinée à la détermination de molécules à utiliser dans le traitement de maladies ou de troubles spécifiques de l'ER $\beta$  ou de maladies ou de troubles associés à un récepteur d'estrogènes selon l'une quelconque des revendications 1 à 8.
15. Utilisation d'un récepteur d'estrogènes selon l'une quelconque des revendications 1 à 8, ou d'une séquence d'ADN selon la revendication 9 ou 10, ou d'une séquence d'acides nucléiques selon la revendication 11 ou 12, dans une analyse destinée à la détermination de molécules à utiliser dans le traitement du cancer de la prostate ou de l'ovaire, de l'adénome prostatique, de maladies du système nerveux central, de l'ostéoporose ou de maladies cardio-vasculaires.
16. Procédé de conception de médicaments comprenant la comparaison de la liaison d'un composé d'essai à l'ER $\alpha$  et à un récepteur d'estrogènes selon l'une quelconque des revendications 1 à 8.
17. Utilisation d'un récepteur d'estrogènes selon l'une quelconque des revendications 1 à 8 dans la recherche d'éventuels effets estrogènes ou d'autres effets hormonaux d'une substance.



FIG. 1

GGAATTCGGGGGAGCTGGCCCAAGGGGAGCGGCTGGTGGCTGGCACTGGCATCCCTAGGC 60  
 ACCCAGSTCTGCAATAAAGTCTGGCAGCCACTGCATGGCTGAGCGACAACCACTGGCTGG 120  
 GAGTCCGGCTCTGTGGCTGAGGAAAGCACTGTCTGCATTAGAGAATGCAAAATAGAGA 180  
 ATGTTAACTGCGCAGTCATTACATCTGAGTCCCATGAGTCTCTGAGAACATAATGTCCAT 240  
 CTGTACCTCTCTCACAAGGAGTTTCTCAGCTGCGACCTCTGAAGACATGGAGATCAA 300  
 AAACACCCGTGAGGCTTAGTTCCTCTCTCTATACTAGAGCAGTCCATCTACCC 360  
 CTGGAGCACGGCCCATCTACATCCCTCTCTCTACGTAGACAACCCCATGAGTATCA 420  
 GCTATGACATTTACAGTCTCTGTGTGATGAACCTACAGTGTTCCTGGCAGCAGTAA 480  
 M T F Y S P A V M N Y S V P G S T S N  
 CTGGACCGTGGGCTGTCCGACTGAGCACAAGCCCAATGTGCTATGGCAACTCTGGG 540  
 L D G G P V R L S T S P N V L W P T S G  
 CACCTGCTCTCTTTAGCGACCCATTGCCAATCATCGCTCTCTATGCAAGCCTCAAAAG 600  
 H L S P L A T H C Q S S L L Y A E P Q K  
 AGTCTTGGTGTGAAGCAAGATCACTAGAGCACACCTTACCTGTAAACAGAGAGACATG 660  
 S F W C E A R S L E H T L P V N E E T L  
 AAGAGGAAGCTTAGTGGGAGCAGTTGTGCCAGCCCTGTACTAGTCCAAACGCAAGAGG 720  
 F F H L S G S S C A S P V T E P N A K F  
 GATGCTCACTCTGCCCCCTGCGAGCGATTATGATCTGGGTATCACTACGGCTTTGG 780  
 D A H F C P V C S D Y A S G Y H V G V W  
 TCATGTGAAGGATGTAAGGCTTTTAAAGAAAGCATTCAAGGACATATGATTATAT 840  
 L E G C K A F E K P S I G G H N D Y I  
 TGTCCAGCCAGCAATCAGTGTACCATAGACAAGAACCGGCGTAAAGCTGCCAGGCTGT 900  
 F P A T N I C T I D E K E E E S C G A S  
 CGACTTGGCAAGTGTATGAAGTAGGAATGGTCAAGTGTGGATCCAGGAGAGAACGGT 960  
 E L F F C V E V G M V K C G S R F E R C  
 GGGTACCTGTATAGTGGGAGGCGAGAGAAGTTCTAGCGAGCAGGTACACTGCCTGAGC 1020  
 G V F I V R R O R S S S E O V H C L S K  
 SCCAAGGAAACGGTGGGCTGCAACCCGGGTGAAGGAGCTACTGCTGAGCAGCTTGAGT 1080  
 A K R N G G H A P R V K E L L L L E T L S  
 CCAGAGCAACTGGTGTCTCACCTCTCTGGAAGCTGAACACCCCAATGTGCTGGTGAAGCCT 1140  
 F E Q L V L T L L E A E P P N V L V S R  
 CCCAGCATGCCCTTACCGAGGCTCCATGATGATGTCCCTCACTAAGCTGGCGGACAG 1200  
 P E M P F T E A S H M H S L T K L A D K  
 GAGCTGGTGCACATGATGGCTGGGCCAAGAAAATCCCTGGCTTTGTGGAGCTCAGCCTG 1260  
 E L V H M I G W A K K I P G F V E L S L  
 TTGGACCAAGTCCGGCTCTTAGAAAGCTGCTGGATGGAGGTGCTAATGGTGGGACTGAT 1320  
 L D L V R L L E S C W N E V L H V G L M  
 TGGTCTCCATCGACCAACCCCGGCAAGCTCATTTTGGCTCCCGACCTCGTTCTGGACGG 1380  
 W F S I D H P G K L I F A P D L V L D R  
 GATGAGGGGAGTGGCTAGAGGGGATCTGGAAATCTTTGACATGCTCTCTGGCGACGACG 1440  
 F E G H C V E G I L E I F D M L L A T T  
 TCAAGCTTCCGTGAGTTAAACCTCCAGCACAAGGAGTATCTCTGTGTGAAGGCCATGAT 1500  
 S F F F R E L K L Q H K E Y L C V K A M I  
 CTCCTCAACTCCAGTATGTACCCCTTGGCTTCTGCAAAACAGGAGGCGAGAAAGTAGCCGG 1560  
 L L H E S M V P L A S A N O E A E S S R  
 AAGGTGACACACCTACTGAACCGCGTGACAGATGCCCTGGTCTGGGTGATTGCGAAGAGT 1620  
 F L T H L L N A V T D A L V W V I A K S  
 GGTATCTCTCTCCAGCAGCAGTCACTCCGACTGGCCAACCTCTGATGCTTCTTTCTAC 1680  
 G C S S C Q C S Y F L A N L L M L L S H  
 CTCAGGSCACATCAGTAAAGGGCATGGAACATCTGCTCAGCATGAAGTGCAAAAATGTS 1740  
 V F H I E N K G M E H L L E K K C K N V  
 GTCCCGGTGTATGACCTGCTGCTGGAGATGCTGAATGCTCACACGCTTCGAGGGTACAAG 1800  
 V P V Y D L L L E M L N A H T L R G Y K  
 TCTTAAATCTCGGGCTCTGAGTGCAGCTCAACAGAGGACAGTAAGAACAAGAGAGCTCT 1860  
 S F I S G S E C S F T E D E K N E E S S  
 CAGAATTTACAGTCTCAGTGTATGGCCAGGCTGAGGCGGACAGACTACAGAGATGGTCAA 1920  
 L H L C S Q  
 AAGTGGAAACATGTACCTAGCATCTGGGGTTCTCTTAGGGCTGCCCTTGGTTACGCACC 1980  
 COTTACTCACACTGCACCTCCAGGAGTCAGGGTGGTGTGTGGCGGTGTTCTCATACC 2040  
 AGGATGTACCAACCGAATGCCAAGTTCTAATCTGTATAGCTTGAAGGCTCTCGGTGTACT 2100  
 TACTTTCTGTCTCTTGGCCACTTGGAAACATCTGAAAGGTTCTGGAACTAAAGGTCAA 2160  
 GTCTGATTGGAAAGGATGTCTTAGTCAGGAAAGGAATATGGCATGTGACACAGCTAT 2220  
 AAGAAATGGACTGTAGGACTGTGTGGCCATAAAATCAACCTTGGATGGCGCTTCTTAGA 2280  
 CCACTTGATTGTAGGATTGAAAACACATTGACAACTCAGCTCATTTCCGATTCTGCTGCT 2340  
 ACCGGCTGTGAGGACTCATTAAATGTCATGGGTATTCTATCAAAAGACGAGAAAGATAGT 2400  
 GCAAGCTTAGATGTACTTGTCTCTCTCCAGACCTTGGGTACATCTCTTAGAGCCTG 2460  
 CTATATGGCTGTCTGAATGTGGTCAATGTGATGGGTAAAGATTAAATCTCTGTGTA 2520  
 TATGGCTCTCTTGAAGTATGTCACTCTCTCTCTCTCTCTCTCTCTCTCTCTCTCTCTCT 2580



FIG. 2A





FIG. 2B

ALIGNMENT OF ER $\beta$  TO OTHER  
ESTROGEN RECEPTORS





## Ligandbinding domain

ELVHMIGWAKKIPG	VELSL	LDQVRL	LESCW	MEVLM	VIL	MMWRS	IDHPGKL	ER8 rat
..N..	RV..	GD.N.H..	H..	CA.L.I..	I..	V..	ME..	ER rat
..N..	RV..	GD.N.H..	H..	CA.L.I..	I..	V..	ME..	ER mouse
..N..	RV..	GD.N.H..	H..	CA.L.I..	I..	V..	ME..	ER human
..N..	RV..	D.T.H..	H..	CA.L.I..	I..	V..	ME..	ER8 rat
IFAPDLVLD	RD	EGK	VEG	ILEIF	DM	LLAT	TTSR	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..	NQ..	MV..	..	..	..	..	..	ER8 rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER rat
..N.L..	NQ..	MV..	..	..	..	..	..	ER mouse
..N.L..	NQ..	MV..	..	..	..	..	..	ER human
..N.L..								

FIG. 2C



## DNA-binding domain

	CPVCS	DYAS	GYHY	GVWS	CEGCKA	FFKR	SIQGH	NDYI	CPAT	NOCTI	DKNRR	KSCOA	CLRK	CYEV	GM
ERA <sup>rat</sup>															
ERA <sup>rat</sup>	A	N							N						
ERA <sup>mouse</sup>	A	N							N						
ERA <sup>human</sup>	A	N							N						

FIG. 2D





FIG. 3A



FIG. 3B




FIG. 3C

FIG. 4A



FIG. 4B




FIG. 4C






SCATCHARD PLOT OF ER-BETA



ER-BETA LIGAND SPECIFICITY











FIG. 7





FIG. 7A

E2 STIMULATED TRANSACTIVATION





DETECTION OF RAT ER EXPRESSION BY RT-PCR.




FIG. 8






FIG. 9



FIG. 10A


HILL PLOT COMPARING hER $\alpha$  AND rER $\beta$ .

FIG. 10B


SCATCHARD PLOT COMPARING hER $\alpha$  AND rER $\beta$ .



FIG.11A




FIG.11B






FIG. 12

Allocation budget of 16, plus 1 for small round checked unlined with 1"AM; 100 resins weight 1140

[illegible]



1 MTFYSPAVMN YSIPSNVTNL EGGPGRQTT PNVLWPTPGH LSPLVVRQL  
 51 SHLYAEPQKS PWCEARSLEH TLPVNRETLK RKVSGNRCAS PVTGPGSKRD  
 101 AHFCAVCSDY ASGYHYGVWS CEGCKAFFKR SIQGHNDYIC PATNQCTIDK  
 151 NRRKSCQACR LRKCYEVGMV KCGSRRERCG YRLVRRQRSA DEQLHCAGKA  
 201 KRSGGHAPRV RELLLDALSP EQVLTLLEA EPPHVLISRP SAPFTEASMM  
 251 MSLTKLADKE LVHMISWAKK IPGFVELSLF DQVRLLESCW MEVLMMGLMW  
 301 RSIDHDPGKLI FAPDLVLRD EGKCEGILE IFDMLLATT RFRELKLQHK  
 351 EYLCVKAMIL LNSSMYPLVT ATQDADSSRK LAHLLNAVTD ALVWVIKSG  
 401 ISSQQQSMRL ANLLMLLSHV RHASNKGMEH LLNMCKKNVV PVDLLLEML  
 451 NAHVLRGCKS SITGSECSA EDSKSKEGSQ NLQSQ\*

FIG. 13A

MAFYSPAVMNYSVPSSTGNLEGGPVRQTASPNVLWPTSGH 40  
 LSPLATHCQSSLLYAEPQKSPWCEARSLEHTLPVNRETLK 80  
 RKLGGSGCASPVTSPTKRDHFCVCSDYASGYHYGVWS 120  
 CEGCKAFFKR SIQGHNDYICPATNQCTIDKNRRKNCQACR 160  
 LRKCYEVGMVKCGSRRERCGYRIVRRQRSASEQVHCLNKA 200  
 KRTSGHTPRVKELLLNSLSPEQLVLTLLAEPPNVLVSRP 240  
 SMPFTEASMMMSLTKLADKELVHMIGWAKKIPGFVELSLL 280  
 DQVRLLESCWMEVLMVGLMWRSIDHDPGKLIFAPDLVLRD 320  
 EGKCEGILEIFDMLLATTARFRELKLQHK EYLCVKAMIL 360  
 LNSSMYHLATASQEAESSRKLTHLLNAVTDALVWVISKSR 400  
 ISSQQQSVRLANLLMLLSHV RHASNKGMEHLLSMKCKNVV 440  
 PVDLLLEMLNAHTLRGYKSSISGSGCCSTEDSKSKEGSQ 480  
 NLQSQ. 486

FIG. 14A



```

1  CTATGACATT CTACAGTCCT GCTGTGATGA ATTACAGCAT TCCCAGCAAT
51  GTCACATACT TCGAAGGTGG GCTGTGTCGG CAGACCACAA GCCCAAATGT
101 GTTGTGGCCA ACACCTGGGC ACCTTTCTCC TTTAGTGGTC CATCGCCAGT
151 TATCACATCT GTATGCGGAA CCTCAAAAGA GTCCCTGGTG TGAAGCAAGA
201 TCGCTAGAAC ACACCTTACC TGTAAACAGA GAGACACTGA AAAGGAAGGT
251 TAGTGGGAAC CGTTGCGCCA GCCCTGTTAC TGGTCCAGGT TCAAAGAGGG
301 ATGCTCACTT CTGCGCTGTC TGCAGCGATT ACGCATCGGG ATATCACTAT
351 GGAGTCTGGT CGTGTGAAGG ATGTAAGGCC TTTTTTAAAA GAAGCATTCA
401 AGGACATAAT GATTATATTT GTCCAGCTAC AAATCAGTGT ACAATCGATA
451 AAAACCGGCG CAAGAGCTGC CAGGCCTGCC GACTTCGGAA GTGTTACGAA
501 GTGGGAATGG TGAAGTGTGG CTCCCGGAGA GAGAGATGTG GGTACCGCCT
551 TGTGCGGAGA CAGAGAAGTG CCGACGAGCA GCTGCACTGT GCCGGCAAGG
601 CCAAGAGAAG TGGCGGCCAC GCGCCCCGAG TCGGGGAGCT GCTGCTGGAC
651 GCCCTGAGCC CCGAGCAGCT AGTGCTCACC CTCCTGGAGG CTGAGCCGCC
701 CCATGTGCTG ATCAGCCGCC CCAGTGCGCC CTTACCCGAG GCCTCCATGA
751 TGATGTCCCT GACCAAGTTG GCCGACAAGG AGTTGGTACA CATGATCAGC
801 TGGGCCAAGA AGATTCCCGG CTTTGTGGAG CTCAGCCTGT TCGACCAAGT
851 GCGGCTCTTG GAGAGCTGTT GGATGGAGGT GTTAATGATG GGGCTGATGT
901 GGCGCTCAAT TGACCACCCC GGCAAGCTCA TCTTTGCTCC AGATCTTGTT
951 CTGGACAGGG ATGAGGGGAA ATGCGTAGAA GGAATTCTGG AAATCTTTGA
1001 CATGCTCCTG GCAACTACTT CAAGGTTTCG AGAGTTAAAA CTCCAACACA
1051 AAGAATATCT CTGTGTCAAG GCCATGATCC TGCTCAATTC CAGTATGTAC
1101 CCTCTGGTCA CAGCGACCCA GGATGCTGAC AGCAGCCGGA AGCTGGCTCA
1151 CTTGCTGAAC GCCGTGACCG ATGCTTTGGT TTGGGTGATT GCCAAGAGCG
1201 GCATCTCCTC CCAGCAGCAA TCCATGCGCC TGGCTAACCT CCTGATGCTC
1251 CTGTCCACG TCAGGCATGC GAGTAACAAG GGCATGGAAC ATCTGCTCAA
1301 CATGAAGTGC AAAAATGTGG TCCCAGTGTA TGACCTGCTG CTGGAGATGC
1351 TGAATGCCCA CGTGCTTCGC GGGTGCAAGT CCTCCATCAC GGGGTCCGAG
1401 TGCAGCCCGG CAGAGGACAG TAAAAGCAAA GAGGGCTCCC AGAACCTACA
1451 GTCTCAGTGA

```

FIG. 13B



ATGCCATTCTAC AGTCTGCTGTG ATGAATACAGT GTTCCAGCAGC ACCGTAACCTG GAAGTGGGCTT 72  
 GTTCGCCAGACT GCAAGCCCAAT GTGCTATGGCCA ACTCTGgACAC CTCCTCTCCTTA GCCACCCACTGC 144  
 CAATCATCGCTT CTCTATGCAGAA CCTCAAAAGAGT CCTTGTGTGAA GCAAGATCACTA GAACACACCTTG 216  
 CCTGTAACAGA GAGACCCCTGAAG AGGAAGCTTGGC GCGAGCGTGTG GCGAGCCCTGTT ACTAGTCCAAgC 288  
 ACCAAGAGGGAT GCTCACTTCTGT GCCGTCTGCACT GATTATGCATCT GGTATCATTTAC GGTGCTCTGGTCC 360  
 TGTGAAGGATGT AAGGCCCTTTT AAAGAAAGCATT CAAGACATAAT GACTATATCTGT CCAGCCACGAAT 432  
 CAGTGTACGATA GACAAGAACCGG CGTAAAACTGC CAGGCCCTGCCGA CTTCCCAAGTGT TAGCAAGTAGGA 504  
 ATGGTCAAGTGT GGATCCAGGAGA GAAAGGTGTGGG TACCGAATAGTA CGAAGACAGAGA AGTCCAGCGAG 576  
 CAGGTGCATTGC CTGAACAAAGCC AAGAGAACCAGT GGGCACACACCC CGGTGAAGGAG CTACTGCTGAAC 648  
 TCTCTGAGTCCC GAGCAGCTGGTG CTACCCCTGCTG GAAGCTGAGCCA CCCAATGTGCTA GTGAGTCTGTCCT 720  
 AGCATGCCCTTC ACCGAGGCTTCC ATGATGATGTCC CTTACGAGCTG GCTGACAAGGAA CTGGTGCACATG 792  
 ATTGGCTGGGC AAGAAATCCCT GGCTTTGTGGAG CTCAGCCTGTG GACCAAGTCCGC CTCCTGGAAAGC 864  
 TGCTGGATGGAG GTGCTGATGGTG GGGCTGATGTGG CGCTCCATCGAC CACCCCGCAAG CTCATCTTTGCT 936  
 CCAGACCTCGET CTGGACAGGGAT GAGGGGAAGTGC GTgAAGGATC CTGGAATCTTT GaCATGCTCCTG 1008  
 GCAGACGACGGCA CGGTTCCGTGAG TTAAACTGCGAG CACAAGATAT CTGTGTGTGAAG GCCATGATTCTC 1080  
 CTCAACTCCAGT ATGTACCACTTG GCTACCCGCAAG GCTAAGGACAG AGTAGCCGGAAG CTGACACACCTA 1152  
 TTGAACGCAGT ACAGATGCCCTG GTCTGGGTGATT TCGAAGAGTAGA ATCTCTTCCcAG CAGCAGTcGTC 1224  
 CGTCTGGCCAAc CTCCTGATGCTT CTTCTCATGTC AGGCACATCAGT AAcAgGGCATG GAACATCTGCTC 1296  
 AGCATGAAGTGC AAAATGTGGTC CCGGTGTACGAC CTGCTGCTGGAG ATGCTGATGCT CACACGCTTCGA 1368  
 GGGTACAAGTCC TCAATCTCGGG TCTGgGTGCTGC TCGACAGAGGAC AGTAAGAGCAAA GAGGGCTCCcAG 1440  
 AACCTCCAGTCT CAGTGA 1458

FIG. 14B



CPD	log IC50 hERu	log IC50 rERu	IC50 (nM) hERu	IC50 (nM) rERu	KI (nM) hERu	KI (nM) rERu	RBA(%) KI hERu	RBA(%) KI rERu	RBA(%) IC50 hERu	RBA(%) IC50 rERu
Dihydrospirodienedione	-6.31	-6.73	485.29	187.11	245.31	163.33	0.027	0.115	0.027	0.115
Testosterone	-5.00	-5.56	10000.00	2187.76	5750.97	1937.70	0.001	0.010	0.001	0.010
Dihydrotestosterone	-6.36	-7.08	436.52	83.95	220.56	73.28	0.030	0.256	0.030	0.256
4-OH-Estradiol	-8.78	-8.67	1.66	2.14	0.95	1.89	6.934	9.892	7.889	10.037
18-Nor testosterone	-5.82	-7.22	1513.56	60.12	765.10	52.48	0.009	0.357	0.009	0.357
5 $\alpha$ -Androstenedione	>-4	>-4	>100000	>100000	>100000	>100000	<0.0002%	<0.0002%	<0.0002%	<0.0002%
Cyprateroneacetate	>-4	>-4	>100000	>100000	>100000	>100000	<0.0002%	<0.0002%	<0.0002%	<0.0002%
8-4 androstene 3,17, dione	>-4	>-4	>100000	>100000	>100000	>100000	<0.0002%	<0.0002%	<0.0002%	<0.0002%
Progesterone	>-4	>-4	>100000	>100000	>100000	>100000	<0.0002%	<0.0002%	<0.0002%	<0.0002%
Coricosterone	>-4	>-4	>100000	>100000	>100000	>100000	<0.0002%	<0.0002%	<0.0002%	<0.0002%
Genistein	-8.35	-9.41	4.47	0.39	2.57	0.34	2.576	54.361	2.931	55.157
$\beta$ -sitosterol	>-4	>-4	>100000	>100000	<0.0002%	<0.0002%	<0.0002%	<0.0002%	<0.0002%	<0.0002%
norethynodrel	-7.53	-7.20	29.51	63.10	14.22	53.09	0.466	0.353	0.444	0.340
norlethindrone	-6.50	-5.89	316.23	1288.25	152.32	1083.89	0.043	0.017	0.041	0.017
$\beta$ -zeaxanthol	-8.80	-9.00	1.29	0.98	0.78	0.87	8.457	21.465	10.162	21.657
O-4 androstien 30,17 $\beta$ -diol	-7.33	-7.64	46.77	22.91	23.39	18.71	0.283	1.001	0.280	0.937
dienestrol	-10.03	-10.46	0.09	0.03	0.05	0.03	140.523	66.418	138.995	618.871
Methoxychlor	-5.45	-8.86	3548.13	109.85	1774.07	89.56	0.004	0.209	0.004	0.196
Bisphenol A	-6.41	-7.07	389.05	42.86	194.52	34.84	0.034	0.538	0.034	0.503
Ecdysterone	>-4	>-4	>100000	>100000	>100000	>100000	<0.0002%	<0.0002%	<0.0002%	<0.0002%
Eudesmide	>-4	>-4	>100000	>100000	>100000	>100000	<0.0002%	<0.0002%	<0.0002%	<0.0002%
Lepidine	>-4	>-4	>100000	>100000	>100000	>100000	<0.0002%	<0.0002%	<0.0002%	<0.0002%
Tachigline	-7.18	-6.63	86.07	234.42	40.13	206.47	0.165	0.091	0.198	0.092
Tachanganidine	-7.87	-8.45	13.49	354.81	8.19	312.50	0.808	0.080	0.971	0.080
Ferulicite	-9.10	-9.58	0.79	0.27	0.40	0.24	16.623	78.091	16.623	78.091
Coumestrol	-9.65	-10.12	0.22	0.08	0.14	0.07	48.665	282.307	58.479	284.839
Naloxidine	-9.32	-9.05	0.48	0.80	0.24	0.78	27.530	23.966	27.530	23.966
16 $\alpha$ -Br-E2	-9.88	-9.87	0.13	0.21	0.07	0.19	100.000	100.000	100.000	100.000
17 $\alpha$ -E2	-9.44	-8.88	0.36	1.32	0.22	1.16	30.006	16.133	38.058	16.278
17 $\beta$ -E2	-9.68	-9.87	0.21	0.13	0.13	0.12	52.145	157.660	62.661	159.074
*19-Ecdysone, 20-Hydroxyecdysone										
RBA-values derived from 16 $\alpha$ -Br-E2 (100%)										

FIG. 15